

# NEWSLETTER GEOBRASIL

## ([www.geobrasil.net](http://www.geobrasil.net))

- **AMAZINGS**

### **VOLCAN SUBMARINO EN LA ANTARTIDA**

Científicos trabajando en las poco hospitalarias aguas de la Península Antártica han encontrado lo que creen es un volcán activo previamente desconocido, situado en el fondo marino. Para confirmarlo, fotografiaron y tomaron muestras de la zona. Los trabajos se han realizado desde el buque oceanográfico Laurence M. Gould, operado por la National Science Foundation, con investigadores de EE.UU. y de Canadá a bordo. El hallazgo del volcán fue casual. Se estaban estudiando las razones por las que una masiva placa de hielo llamada Larsen B se derrumbó y cayó al agua hace varios años (los científicos quieren saber si este fenómeno fue una situación única o si forma parte de un ciclo que se extiende a lo largo de cientos de miles de años). Ya hacía tiempo, además, que los marineros de diversas expediciones habían informado de que las aguas en la zona tenían un color peculiar. Después de la exploración preliminar, Eugene Domack, el científico jefe de la última expedición, ha manifestado que la información obtenida permite afirmar que el volcán se eleva unos 700 metros por encima del fondo marino, hasta unos 275 metros por debajo del nivel del mar. Según las estimaciones, el cono volcánico poseería al menos 1,5 kilómetros cúbicos de roca volcánica. Sirva como comparación el Monte Erebus, un volcán activo situado en la isla de Ross, cerca de la McMurdo Station, que tienen unos 3.800 metros de altitud sobre el nivel del mar. El mayor volcán de la Tierra, el Mauna Loa hawaiano, se eleva aproximadamente 4.100 metros, mientras que el más pequeño, el Cuexcomate, de México, mide apenas 13 metros de alto. Hasta ahora no se tenía constancia de la existencia de volcanes activos en la región llamada Antarctic Sound, en la punta más al norte de la Antártida. De hecho, el pico recién descubierto está al norte del límite debajo del cual sí se sabía que existía actividad. Los mapas levantados por el sónar en enero de 2002 dieron las primeras pistas de la existencia del volcán. Ponían de manifiesto una gran estructura concéntrica y simétrica que no había sido afectada por el avance o retroceso de los glaciares. Ello hace sospechar que se trata de un ejemplar joven, geológicamente hablando. Fue en abril de 2004 cuando los científicos pudieron regresar para obtener más evidencias. Así, se emplearon cámaras de video, sondas de temperatura y sistemas para dragar rocas. Las imágenes muestran que la superficie está altamente colonizada por organismos, excepto en algunas zonas donde la negra roca indica que la lava ha fluido recientemente por ahí. Las operaciones de dragado, además, permitieron capturar numerosas muestras de roca basáltica. Las sondas de temperatura, por su parte, detectaron puntos calientes cerca de los lugares en los que la roca es más joven. Información adicional en: <http://www.amazings.com/ciencia/noticias/280504c.html>

- **NATURE**

Electric polarization reversal and memory in a multiferroic material induced by magnetic fields 392

N. HUR, S. PARK, P. A. SHARMA, J. S. AHN, S. GUHA & S-W. CHEONG

doi: 10.1038/nature02572

[First paragraph](#) | [Full Text](#)

Evidence from massive siderite beds for a CO<sub>2</sub>-rich atmosphere before ~ 1.8 billion years ago 395

HIROSHI OHMOTO, YUMIKO WATANABE & KAZUMASA KUMAZAWA

doi: 10.1038/nature02573  
[First paragraph](#) | [Full Text](#)

- **SCIENCE**

M Andersson, RT Ottesen, and T Volden Building materials as a source of PCB pollution in Bergen, Norway. *Sci Total Environ* 5 Jun 2004 325(1-3): p. 139.  
<http://highwire.stanford.edu/cgi/medline/pmid;15144784>

May Nyman, Francois Bonhomme, Todd M. Alam, John B. Parise, and Gavin M. B. Vaughan [SiNb<sub>12</sub>O<sub>40</sub>]<sup>16-</sup> and [GeNb<sub>12</sub>O<sub>40</sub>]<sup>16-</sup>: Highly Charged Keggin Ions with Sticky Surfaces. *Angew Chem Int Ed Engl* 17 May 2004 43(21): p. 2787.  
<http://highwire.stanford.edu/cgi/medline/pmid;15150750>

C Franzen, R Killian, and H Biester Natural mercury enrichment in a minerogenic fen-evaluation of sources and processes. *J Environ Monit* 1 May 2004 6(5): p. 466.  
<http://highwire.stanford.edu/cgi/medline/pmid;15152316>

W Shotyk and M Krachler Atmospheric deposition of silver and thallium since 12 370 (14)C years BP recorded by a Swiss peat bog profile, and comparison with lead and cadmium. *J Environ Monit* 1 May 2004 6(5): p. 427.  
<http://highwire.stanford.edu/cgi/medline/pmid;15152311>

M Krachler and W Shotyk Natural and anthropogenic enrichments of molybdenum, thorium, and uranium in a complete peat bog profile, Jura Mountains, Switzerland. *J Environ Monit* 1 May 2004 6(5): p. 418.  
<http://highwire.stanford.edu/cgi/medline/pmid;15152310>

G Le Roux, D Weiss, J Grattan, N Givelet, M Krachler, A Cheburkin, N Rausch, B Kober, and W Shotyk Identifying the sources and timing of ancient and medieval atmospheric lead pollution in England using a peat profile from Lindow bog, Manchester. *J Environ Monit* 1 May 2004 6(5): p. 502.  
<http://highwire.stanford.edu/cgi/medline/pmid;15152320>

N Givelet, G Le Roux, A Cheburkin, B Chen, J Frank, ME Goodsite, H Kempter, M Krachler, T Noernberg, N Rausch, S Rheinberger, F Roos-Barraclough, A Sapkota, C Scholz, and W Shotyk Suggested protocol for collecting, handling and preparing peat cores and peat samples for physical, chemical, mineralogical and isotopic analyses. *J Environ Monit* 1 May 2004 6(5): p. 481.  
<http://highwire.stanford.edu/cgi/medline/pmid;15152318>

Z Wu, J Wang, M Zhuo, and H Wan [Spatial distribution and landscape ecological impact degree assessment of quarry in Zhuhai City] *Ying Yong Sheng Tai Xue Bao* 1 Feb 2004 15(2): p. 283. <http://highwire.stanford.edu/cgi/medline/pmid;15146640>

Impact of the Capillary Fringe on Local Flow, Chemical Migration, and Microbiology Brian Berkowitz, Stephen E. Silliman, and Andrea M. Dunn *Vadose Zone J* 1 May 2004; 3(2): p. 534-548 <http://vzj.scijournals.org/cgi/content/abstract/3/2/534?ct>

Modeling Boron Adsorption Isotherms and Envelopes Using the Constant Capacitance Model Sabine Goldberg *Vadose Zone J* 1 May 2004; 3(2): p. 676-680  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/676?ct>

Capillary Pressure-Saturation Relations for Saprolite: Scaling With and Without Correction for Column Height E. Perfect, L. D. McKay, S. C. Cropper, S. G. Driese, G. Kammerer,

and J. H. Dane Vadose Zone J 1 May 2004; 3(2): p. 493-501  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/493?ct>

Comparison of Hanford Colloids and Kaolinite Transport in Porous Media Jie Zhuang, Yan Jin, and Markus Flury Vadose Zone J 1 May 2004; 3(2): p. 395-402  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/395?ct>

Flint mining in prehistory recorded by in situ-produced cosmogenic  $^{10}\text{Be}$  G. Verri, R. Barkai, C. Bordeanu, A. Gopher, M. Hass, A. Kaufman, P. Kubik, E. Montanari, M. Paul, A. Ronen, S. Weiner, and E. Boaretto Proc. Natl. Acad. Sci. USA 25 May 2004; 101(21): p. 7880-7884  
<http://www.pnas.org/cgi/content/abstract/101/21/7880?ct>

Atomic force microscopy of Precambrian microscopic fossils Andre Kempe, J. William Schopf, Wladyslaw Altermann, Anatoliy B. Kudryavtsev, and Wolfgang M. Heckl Proc. Natl. Acad. Sci. USA published 25 May 2004, 10.1073/pnas.0403146101  
<http://www.pnas.org/cgi/content/abstract/0403146101v1?ct>

Impact of the Capillary Fringe on Local Flow, Chemical Migration, and Microbiology Brian Berkowitz, Stephen E. Silliman, and Andrea M. Dunn Vadose Zone J 1 May 2004; 3(2): p. 534-548  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/534?ct>

Modeling Boron Adsorption Isotherms and Envelopes Using the Constant Capacitance Model Sabine Goldberg Vadose Zone J 1 May 2004; 3(2): p. 676-680  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/676?ct>

Capillary Pressure-Saturation Relations for Saprolite: Scaling With and Without Correction for Column Height E. Perfect, L. D. McKay, S. C. Cropper, S. G. Driese, G. Kammerer, and J. H. Dane Vadose Zone J 1 May 2004; 3(2): p. 493-501  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/493?ct>

Comparison of Hanford Colloids and Kaolinite Transport in Porous Media Jie Zhuang, Yan Jin, and Markus Flury Vadose Zone J 1 May 2004; 3(2): p. 395-402  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/395?ct>

Soil Mineralogy with Environmental Applications. Robert C. Graham Vadose Zone J 1 May 2004; 3(2): p. 724  
<http://vzj.scijournals.org/cgi/content/full/3/2/724?ct>

Water-Dispersible Colloids: Effects of Measurement Method, Clay Content, Initial Soil Matric Potential, and Wetting Rate Charlotte Kjaergaard, Lis W. de Jonge, Per Moldrup, and Per Schjonning Vadose Zone J 1 May 2004; 3(2): p. 403-412  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/403?ct>

Anion Exchange in Saprolite K. A. McVay, D. E. Radcliffe, L. T. West, and M. L. Cabrera Vadose Zone J 1 May 2004; 3(2): p. 668-675  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/668?ct>

Rainfall-Induced Soil Surface Sealing: A Critical Review of Observations, Conceptual Models, and Solutions S. Assouline Vadose Zone J 1 May 2004; 3(2): p. 570-591  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/570?ct>

Measurement of the Solid Dielectric Permittivity of Clay Minerals and Granular Samples Using a Time Domain Reflectometry Immersion Method D. A. Robinson Vadose Zone J 1 May 2004; 3(2): p. 705-713  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/705?ct>

Geochemical and Hydrological Reactivity of Heavy Metals in Soils. Molly N. Pace Vadose Zone J 1 May 2004; 3(2): p. 733  
<http://vzj.scijournals.org/cgi/content/full/3/2/733?ct>

Faulting and Fracturing of Nonwelded Bishop Tuff, Eastern California: Deformation Mechanisms in Very Porous Materials in the Vadose Zone James P. Evans and Kelly Keighley Bradbury *Vadose Zone J* 1 May 2004; 3(2): p. 602-623  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/602?ct>

Cyanide Leaching from Soil Developed from Coking Plant Purifier Waste as Influenced by Citrate Tim Mansfeldt, Heike Leyer, Kurt Barmettler, and Ruben Kretzschmar *Vadose Zone J* 1 May 2004; 3(2): p. 471-479  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/471?ct>

Particle Leaching and Particle-Facilitated Transport of Phosphorus at Field Scale L. W. de Jonge, P. Moldrup, G. H. Rubaek, K. Schelde, and J. Djurhuus *Vadose Zone J* 1 May 2004; 3(2): p. 462-470  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/462?ct>

Colloid Mobilization and Transport in Undisturbed Soil Columns. II. The Role of Colloid Dispersibility and Preferential Flow Charlotte Kjaergaard, Per Moldrup, Lis W. de Jonge, and Ole H. Jacobsen *Vadose Zone J* 1 May 2004; 3(2): p. 424-433  
<http://vzj.scijournals.org/cgi/content/abstract/3/2/424?ct>

Reflecting on Earth's Climate Science 28 May 2004; 304(5675): p. 1207f  
<http://www.sciencemag.org/cgi/content/summary/304/5675/1207f?ct>

DEVELOPMENTAL BIOLOGY: Enhanced: The Ups and Downs of a Sea Anemone Peter Holland *Science* 28 May 2004; 304(5675): p. 1255-1256  
<http://www.sciencemag.org/cgi/content/summary/304/5675/1255?ct>

PALEOCEANOGRAPHY: Assessing Sediment Age H. Jesse Smith *Science* 28 May 2004; 304(5675): p. 1215b  
<http://www.sciencemag.org/cgi/content/summary/304/5675/1215b?ct>

High Pore Fluid Pressure May Cause Silent Slip in the Nankai Trough Shuichi Kodaira, Takashi Iidaka, Aitaro Kato, Jin-Oh Park, Takaya Iwasaki, and Yoshiyuki Kaneda *Science* 28 May 2004; 304(5675): p. 1295-1298  
<http://www.sciencemag.org/cgi/content/abstract/304/5675/1295?ct>

Fluid Source for Seismic Slip *Science* 28 May 2004; 304(5675): p. 1207f  
<http://www.sciencemag.org/cgi/content/summary/304/5675/1207f?ct>

## • EARTH PAGES

Web resources

### **Impacts' effects**

Algorithms that model the physical effects of extraterrestrial impacts from the Lunar and Planetary Laboratory of the University of Arizona, headed by Jay Melosh, have been assembled into a handy on-line calculator, with notes on the processes involved. If you want to find out if you will be fried, buried or blown to smithereens (probably all three if our luck is really out), and the chances of being harmed by alien lumps of rock or ice, you can find the calculator at <http://www.lpl.arizona.edu/impacteffects/>. It is not recommended for estate agents, because, unlike many other disastrous events, impacts can be anticipated anywhere.

Anthropology and geoarchaeology

### **Early humans of Beijing**

One of the most remarkable achievements of early humans (*Homo ergaster* aka *H. erectus*) was not their tools, but their migration out of Africa around 1.8 Ma, to reach as far as Indonesia and China. There is no evidence for that feat having occurred again until fully modern humans arrived in east Asia about 70 ka ago. The toolkit of Asian "Action

Man" is unimpressive, in the sense that it resembles the slightly reshaped broken pebbles of the Oldowan culture, that first appears in the African archaeological record about 2.4 Ma ago. Development in Africa of the enigmatic and beautiful bi-face or Acheulean axe was after the first Asians had departed, around 1.5 Ma. So what were these early wanderers like; what did they want? The decade-long work in China by Noel Boaz, an anatomist from the Ross School of Medicine in New Jersey and anthropologist Russell Ciochon of the University of Iowa will soon appear in their book *Dragon Bone Hill, an Ice-Age Saga of Homo Erectus* (Oxford University Press), which they preview in the 17 April 2004 issue of *New Scientist* (p. 32-35). Boaz and Ciochon have worked mainly in Zhoukoudian near Beijing, a major resource for human remains whose different levels extend back to about 800 thousand years. Another site in China, Longouppo, contains disputed remains as old as 1.8 Ma, as are Dubois' famous discoveries of the type specimens of *H. erectus* by the Solo River in Java. From the time when Zhoukoudian became famous among Chinese apothecaries as a source of "dragon's bones" (a mixture of human and other animal remains) there has always been an air of myth about the findings there – a permanent dwelling for hundreds of thousand years, protected from glacial temperature falls by the consistent use of fire. In essence, the publicised view is that "Peking Man" led a cosy hearthside existence for a very long time indeed. Boaz and Ciochon tell a different, and more mundane story. Most bones in the deposit are those of a great variety of other animals, with disproportionately few of human origin, and those are highly fragmented. The dominant species is a giant hyena, and many of the bones, including humans, are well gnawed, which is what hyenas do especially well. There are occasional signs of human occupation and use of fire. The human remains are encased in layered carbonate flowstone,. Records of fluctuating  $d^{18}O$  from that matrix, matched against the global time series of climate change, show that occupation was only during interglacials - the site was abandoned or unvisited during the depth of glacial periods. Some animal bones show cut marks made by stone tools, and it is more likely that *H. erectus* raided to get remnants of other beasts' kills, perhaps using fire, rather than being top of the predatory order. The great surprise throughout Asia is the complete lack of development of stone tools from the primitive culture that arrived there, until as late as 20 to 30 thousand years ago, when Asian *H. erectus* vanished. Apart from the stunning breakthrough to the bi-face axe, African erects also had a million-year long cultural stasis – resting on laurels with a vengeance. Finally, from a number of skulls at Zhoukodian, Boaz and Ciochon have shown signs of trauma. These are depression fractures, probably not necessarily fatal, but indicate sharp blows to the head with blunt instruments. Their interpretation is that the Chinese erects settled disputes by bashing heads; so that aspect of culture has not changed a lot since. Their story is not "politically correct", but with publication of their book, other palaeoanthropologists can judge it on the basis of the evidence from Dragon Bone Hill.

#### **Faster development of Neanderthals**

Go to any horse sale and you will see bidders closely studying the teeth of their prospective purchases; the origin of the saying, "Never look a gift horse in the mouth". Teeth show growth ridges, and in grazing animals they are prominent, so that it is possible to judge the age of a horse easily and accurately. Human teeth are different only in the less obvious signs of growth. Microscopic examination reveals such records, down to the daily level, although the most prominent features are curious disturbances in their deposition that form approximately weekly. They appear as ridges on the crowns of teeth. The variable spacing of these *perikymata* provides a record of the pace at which adult teeth develop. In modern humans the spacing becomes very much closer in the later growth history (towards the tooth's cutting edge) than in its early stages, and reflects the slow development to full adult dentition. In a painstaking study of hundreds of teeth from Cro Magnon and Neanderthal teeth, Fernando Rozzi of the University of Paris and José Bermudez de Castro of the Spanish National Museum of Natural Sciences have discovered an odd difference in the development rates of Neanderthals (Rozzi, F.V.R & Bermudez de Castro, J.M. 2004. Surprisingly rapid growth in Neanderthals. *Nature*, v. **428**, p. 936-939). The late *perikymata* of Neanderthals are more widely spaced than in Cro Magnon and modern humans, strongly suggesting that Neanderthals

developed to adulthood by about the age of 15, three to five years earlier than us and our immediate ancestors. As well as confirming that they are a separate species, the results suggest that Neanderthals, while acquiring brains as large, and in some cases even larger than ours, had evolved more rapid maturation and probably a genetically determined shorter adult life. This would have had some effect on transfer of culture, which in human societies is often the most important value of elderly folk. The fewer samples of teeth of earlier human species (*H. heidelbergensis* and *H. antecessor*) reveal an even greater surprise. They are more like modern human teeth (albeit with signs of somewhat faster growth), which suggests that evolution of the Neanderthals involved a regression. The authors suggest that the combination of a backward step to faster development with rapid brain growth to large size might reflect a very-high calorie diet together with adverse environmental conditions.

Geobiology, palaeontology, and evolution

#### **Devonian broad-shouldered fish**

How, when and under what circumstances vertebrates got limbs to take them charging across the forested land of the late Palaeozoic form a central issue in our own evolution, as well as that of the other four-footed land animals. By negative analogy with the functional though rather rudimentary enlarged fins of various modern fish that flop from pond to pond during dry seasons, many vertebrate palaeontologists have considered limbs as evolutionary adaptations in air-breathing fish once they made this a habit. As so often, the fossil record has not given up enough evidence for that to be certain. Well, an upper foreleg bone (*humerus*) has turned up in Late Devonian rocks from Pennsylvania at a time and in a context that strongly suggests it was carried by a fish (Shubin, N.H. *et al.* 2004. The early evolution of the tetrapod humerus. *Science*, v. **304**, p. 90-93). While not able to ride a bicycle, the advanced fish probably used what became limbs to hold itself motionless while lying in ambush for its prey. That would provide a plausible point of departure from which walking might develop.

#### **Early biomarkers in South African pillow lavas**

It is now established that various kinds of bacteria infest rocks down to depths of 2 km or more, one particularly favourable habitat being in sea-floor basalts through which hydrothermal fluids travel. Although the majority probably inhabits cracks and joints, some seem to work actively to corrode rock, especially volcanic glass, thereby obtaining mineral nutrients. Signs of this microbial corrosion in modern volcanic glasses are radiating tubes on a scale of a few micrometres, that show up in micrographs, and many may have been overlooked by petrographers in all kinds of rock. That they are definitely formed by organic activity is demonstrated by the presence of nucleic acids, carbon and nitrogen in the tubules. Carbon isotopes from them show the strong depletion in  $^{13}\text{C}$  that is the hallmark of organic fractionation of natural carbon. A team of geoscientists, from Norway, Canada and the USA, who have steadily accumulated evidence for biological rotting in modern oceanic basalts, turned their focus to the oldest, well-preserved pillow lavas in the 3.5 billion-year old Barberton greenstone belt of north-eastern South Africa (Furnes, H. *et al.* 2004. Early life recorded in Archean pillow lavas. *Science*, v. **304**, p. 578-581). Virtually identical microtubules seem common in them too, particularly in hydrated glasses that are now tinged with the low-grade metamorphic mineral chlorite. Indeed, chlorite seems to have grown preferentially from clusters of the holes, which suggests that they formed before metamorphism of the basalts. Micro-geochemical studies confirm the presence of hydrocarbons with low  $\delta^{13}\text{C}$ . The bulk of the tubules occur in the inter-pillow debris, that probably formed as glassy rinds as magma protruded on the Archaean sea floor. As well as adding to evidence for ancient terrestrial life, the find has inevitably opened up the search for such signs in meteorites reckoned to have come from Mars. In two, olivine grains show similar structures, although why the olivine hadn't broken down in the presence of water that is essential for life makes such observations worth taking with a pinch of salt. A number of studies have stymied claims for early bacterial fossils (see *Artificial Archaean "fossils"* and *Doubt cast on earliest bacterial fossils*, April 2002 and December 2003 issues of *EPN*) and inorganic processes conceivably might create structures that can be mistaken for ones formed by biological action. The Fischer-Tropsch process is capable of producing hydrocarbons, and produces



depletion in  $^{13}\text{C}$  abiogenically. In the on-line April edition of Science Express (<http://www.scienceexpress.org>) experiments are reported that highlight the possible influence of chromium-bearing mineral catalysts in hydrothermal generation of hydrocarbons from inorganic carbon dioxide (Foustoukos, D.I. & Seyfried, W.E. 2004. Hydrocarbons in hydrothermal vent fluids: the role of chrome-bearing catalysts. *Science Express*, April 2004). The Barberton greenstone belt is well known for ultramafic lavas rich in chromium, as are most early volcanic sequences.

**See also:** Kerr, R.A. 2004. New biomarker proposed for earliest life on Earth. *Science*, v. **304**, p. 503.

Geochemistry, mineralogy, petrology and volcanology

**And now....molybdenum isotopes! Ocean anoxia in the Proterozoic**

"Everyone knows" that free atmospheric oxygen appeared about 2300 million years ago, thanks to the waste products of blue-green bacterial photosynthesis. At least the land surface became an oxidising environment and a progressively redder place, as Fe-2 was oxidised to Fe-3 which forms insoluble oxides and hydroxides. Paradoxically, the shallow sea floor of earlier times was redder than anything since, because of exactly the same oxygen-containing, ferric minerals. It hosted the largest build-up of any metal concentration in Earth's history; the banded iron formations (BIFs) that have for a century or more been the source of industrial iron. A simple, and probably accurate explanation for BIFs is that iron dissolved in ocean water that lacked oxygen as Fe-2, and was supplied by sea-floor volcanism. Once blue-green bacteria began pumping out oxygen, an oxidising reaction dumped both elements as slimy red sediment where the two met. Dissolved iron consumed oxygen – just as well, because to most prokaryote life it is a poison – yet as oxygen productivity rose (and perhaps sea-floor spreading slowed) dissolved iron was increasingly removed by oxidation from sea water. The tipping point, when air contained oxygen and sea water became starved of iron (a vital micronutrient for phytoplankton) is difficult to address since the two chemical environments are so different and interact in complicated ways. BIFs continued to form for about half a billion years after the first sign of atmospheric oxygen, then they disappear from the geological record at 1800 Ma ago. There were minor reappearances in the Neoproterozoic, at the time of "Snowball Earth" events, and that is a fascinating topic in its own right. Clearly, there was a long period of transition to what we can regard as a thoroughly modern world. Studies that use sulphur isotopes suggest that in the Mesoproterozoic the upper ocean was oxygenated while bottom waters were perpetually akin to those of the Black Sea today. Conditions in them may have been highly conducive to burial of dead organic matter – rapid drawdown of atmospheric  $\text{CO}_2$ , but allowing the massive production of methane by anaerobic bacteria. Methane is a far more potent greenhouse gas than carbon dioxide, so controls over climate may have been very different from today's. Molybdenum offers an independent and potentially useful means of testing hypotheses about ocean chemistry. It enters the sea in river water, which in post 2300 Ma times would have been oxygenated, allowing the formation of the soluble and very stable molybdate ion. In anoxic ocean floor conditions, bacteria that generate hydrogen sulphide remove molybdenum as the sulphide, which is why modern Mo concentrations remain stable – it ends up in a very small percentage of ocean floor sediments. The stable isotopes of molybdenum ( $^{97}\text{Mo}$  and  $^{95}\text{Mo}$ ) fractionate during precipitation of the element, the heavier one being preferentially removed during sulphide precipitation, to give high  $^{97}\text{Mo}/^{95}\text{Mo}$  ratios in sediments. The opposite seems to occur if precipitation is in the oxide form, as in sea-floor manganese nodules. Geochemists from the Universities of Rochester and Missouri, USA have compared Mo isotopes from apparently anoxic Mesoproterozoic sediments with those in modern euxinic basins (Arnold, G.L. *et al.* 2004. Molybdenum isotope evidence for widespread anoxia in mid-Proterozoic oceans. *Science* v. **304**, p. 87-90). The Precambrian results are isotopically much lighter than modern ones, suggesting that  $^{97}\text{Mo}$  did not become enriched in seawater as a result of oxide precipitation in the equivalent of modern manganese nodules. They estimate that 10 times more of the ocean floor was anoxic than today or since about 1300 Ma ago. So far no comparable work has been done of the extremely abundant black shales and schists of the Neoproterozoic, that link with "Snowball Earth" events. Whether or not "modern"

redox conditions emerged 1300 Ma ago, with probably a big impact on climate controls, the oddest time climatically was between about 750 and 600 Ma ago. Not only were there several dramatic coolings and warmings, but the main indicator of organic carbon burial,  $d^{13}C$ , went haywire. As did the BIFs, did ocean anoxic conditions once more get footholds. Molybdenum isotope data seem likely to shed some light on those strange times.

Geomorphology

### **River incision and anticlines**

In many areas of active deformation, landforms that suggest that uplift and river down-cutting keep pace are very common. Stream courses cross zones of uplift, rather than being diverted or ponded up to form lakes. Traditionally, geomorphologists have described such drainages as "antecedent", i.e. rivers that were present before uplift began. They can be seen on all scales up to examples such as the Indus and Brahmaputra rivers that carve their way across the actively rising Himalaya. The most common are anticlines through which streams flow in canyons perpendicular to the fold axes. A curious and common feature is that the canyons are not haphazard, but often cut the fold where its amplitude is greatest and its axis plunges away from the site of incision. The stupendous rates at which crustal rocks are eroded and transported away in the courses of the Indus and Brahmaputra, and in lesser drainages on the flanks of major extensional orogens, such as the Red Sea, clearly removes load from the crust. Consequently there is an isostatic component to the uplift involved in the two cases at a grand scale. Peter Molnar and Phillip England suggested an erosional role in large-scale uplift over a decade ago. Intervening ridges rise higher than they would if erosion was slower or non-existent. In major rift systems, the highest peaks are often within the escarpments rather than at the lip of uplift, sometimes more than 500 m higher. Bearing this well-known process in mind, Guy Simpson of ETH Zurich, has sought evidence that it functions on much smaller scales (Simpson, G. 2004. Role of river incision in enhancing deformation. *Geology*, v. **32**, p. 341-344). That comes from the surprising symmetry of doubly plunging anticlines that are cut by rivers at their highest point. His modelling suggests that the phenomenon can occur when the crust deforms plastically, allowing isostatic response to erosion on even minor scales during compression. When deformation is by brittle means, any uplift of rigid crust is flexural and has long wavelengths, so that rivers bear no relation to local structures

Planetary, extraterrestrial geology, and meteoritics

### **Water on Mars; almost official**

Two lines of evidence from the current robotic explorations of Mars add to less tenuous ones that the planet is really wet – icy to be precise. One is mineralogical. Spectroscopy of the surface being slowly trundled across by a NASA rover, shows abundant signs of the hydrated, iron-potassium sulphate jarosite, which probably can only form under wet conditions. When it was precipitated is not known with certainty, but it occurs in layered sediments that contain structures that clearly point to transport in and deposition from surface water. The time when liquid water could exist at the surface probably goes back to the earliest events on Mars, tied to the famous canyons and more recently discovered dendritic drainage patterns. The other evidence stems from even more remote sensing, that captures short-wavelength infrared radiation emitted by the Sun and reflected from the Martian surface. Ices of water and carbon dioxide have distinct and unique reflected spectra, because of the different ways in which they absorb a small proportion of solar radiation. Results from the OMEGA instrument aboard the European Space Agency's Mars Express satellite show that the south polar region contains as much as 15% water ice mixed with solid  $CO_2$  (Bibring, J-P *et al.* 2004. Perennial water ice identified in the south polar cap of Mars. *Nature*, v. **428**, p. 627-630).

Sedimentology and stratigraphy

### **Magnetic polarity reversals**

The Earth's magnetic field is changing all the time, in its intensity, direction and, now and again, its polarity. It's the last that proved the key to sea-floor spreading and plate tectonics, though ocean-floor magnetic "stripes", and which has become a key stratigraphic tool for correlation and approximate dating. Along with palaeomagnetic pole



determinations, that are vital to continental reconstructions, the whole field still remains largely empirical. Although widely agreed to be connected to changes in motions in the core, exactly what happens during reversals of geomagnetic polarity remains enigmatic, despite 40 years having passed since they were first recognised. There is no doubt that they are quick events, but to judge their pace and what happens to field strength and direction during a "flip" requires high quality data that is well-calibrated to time. Most early work focussed on magnetisation in igneous rocks, where the signal is strong. Minerals such as igneous magnetite acquire a permanent magnetisation once they cool below their Curie temperature, but since accurate radiometric dating gives an age, not a range of ages, it might seem that all that is possible with lavas and intrusions is to obtain a series of points. Fine for a time series, but useless for the details of reversals. However, by modelling the cooling history of an igneous body, it is possible to calibrate different levels within it to time. With careful choice, it has proved possible to find flows in flood basalt sequences that include the brief progress of a reversal. The results seem very odd, the pole itself seeming to migrate rather than jump from north to south, and gross changes in intensity over a short time. Improved instrumentation allows a shift from strongly magnetic basalts, to sediments that preserve much weaker signals. These are due to the alignment with the field of magnetic grains as they slowly settle. Marine sediment cores can now be magnetically characterised – the principle behind magnetostratigraphy. For geomagnetists the most recent reversals have proved especially instructive, when the sedimentary record is analysed (Clement, B.M. 2004. Dependence of the duration of geomagnetic polarity reversals on site latitude. *Nature*, v. **428**, p. 637-640). On average, the last four "flips" took about 7000 years to complete by migration of the magnetic poles. Yet there is an oddity in the detail. Sites at low latitude show significantly shorter periods (down to 2000 years) than those at high latitude (as much as 10000 years). Clement's explanation for the difference is the persistence of the lower intensity non-dipole field, which might suggest different core processes or a single process with several components that evolve at different rates.

#### **Sulphur cycling and sea-level change**

Sulphur is one the major prerequisites for life after carbon, hydrogen, oxygen and nitrogen, and the bulk of it is supplied by sulphate ions. After chlorine, the  $\text{SO}_4^{2-}$  ion is the most abundant anion in the oceans. Not very much is added annually by river drainage, and although anaerobic bacteria remove some by reducing it to hydrogen sulphide so that it is removed from solution as a result of precipitation of insoluble iron sulphide, the sulphur cycle has been considered to be the most sluggish of all the major geochemical rhythms at the Earth's surface. Because iron sulphide is highly reactive in oxidising conditions, should marine sulphide-rich sediments become exposed at the surface their oxidation to sulphuric acid and iron hydroxide would rapidly add sulphate ions to seawater. Studies of sulphur isotopes seem to suggest that this is not very important however. Through sulphate-sulphide reducing bacteria, sulphur is implicated in the carbon cycle because of its sheer abundance, not so much from the encouragement and burial of the bacteria, but because they induce the highly reducing conditions that help a larger proportion of dead organic matter to remain unoxidised and become buried. In a roundabout way, sulphur has a role in climate controls. In fact, two roles. Sulphate ions affect the alkalinity of seawater, and on that depends the oceans' ability to dissolve  $\text{CO}_2$  from the atmosphere. The big question is, "Does the sulphate content of seawater ever change fast enough to have some impact on climate in the short term?". Most studies of the S-cycle have focused on sulphur isotopes, so a new twist is bound to be interesting. Alexandra Turchyn and Daniel Schrag of Harvard University looked instead at the isotopes of oxygen within barium sulphate contained within seafloor sediments since the Late Miocene (about 10 Ma ago) (Turchyn, A.V. & Schrag, D.P. 2004. Oxygen isotope constraints on the sulfur cycle over the past 10 million years. *Science*, v. **303**, p. 2004-2007). Up until 6 Ma, the barite  $d^{18}\text{O}$  (measured against mean ocean water values) stayed constant at about 9.5‰, and then rose to around 12.5‰ by 3.5 Ma. Through the Late Pliocene and Pleistocene, the period of repeated glacial-interglacial cycles, it fell dramatically to its present level of 7.9‰. In that later period, the average  $d^{16}\text{O}$  of deep water foraminifera rose significantly. The decline in "heavy" oxygen in marine sulphates

can be linked to increased exposure of pyrite-bearing marine sediments during glacial sea-level falls when "light" atmospheric oxygen enters the sulphate ions that are produced. Modelling suggests sulphate ions in seawater increased by as much as 20% during the Great Ice Age. Whether that had an influence on the oceans' take-up of carbon dioxide from the atmosphere in the last 3 Ma is yet to be evaluated. However, Turchyn and Schrag's detection of a short term shift in the sulphur cycle, and attributing it to falling sea level, may allow a new approach to global sea-level change, which has mainly been deduced from features in stratigraphy.

**See also:** Derry, L.A. & Murray, R.W. 2004. Continental margins and the sulfur cycle. *Science*, v. **303**, p. 1981-1982

- **IAPC**

*Geochemistry International*

Vol. 42, No. 5, 2004

Simultaneous English language translation of the journal is available from MAIK "Nauka / Interperiodica" (Russia).

*Geochemistry International* ISSN 0016-7029.

Composition and Sources of Magmas in Medvezh'ya Caldera (Iturup Island, Southern Kuriles)

from a Study of Melt Inclusions

V. I. Kovalenko, V. B. Naumov, M. L. Tolstykh,  
G. M. Tsareva, and N. N. Kononkova p. 393 [abstract](#)

Geochemistry of a Neutrally Buoyant Plume above the Rainbow Hydrothermal Field and Related Fluxes of Sedimentary Material

V. N. Lukashin, D. L. Aleinik, A. B. Isaeva, A. Yu. Lein, and G. M. Vinogradov p. 414 [abstract](#)

Pore Solutions in the Rocks of the Schlema Uranium Deposit, Erzgebirge, and Their Role in Ore Formation

Vikt. L. Barsukov p. 428 [abstract](#)

Genesis and Composition of Borates of the Ludwigite–Vonsenite Series in Magnesian Skarns

of the Urals, Siberia, and the Russian Far East

S. M. Aleksandrov and M. A. Troneva p. 449 [abstract](#)

Genesis of Stratiform Pb–Zn Deposits in Carbonate Rocks: A Hydrogeochemical Approach

B. N. Ryzhenko and S. R. Krainov p. 465 [abstract](#)

Short Communications

Isotopic Geochemical Parameters of the Formation of Schorl–Dravite Tourmaline from Hydrothermal Deposits in the Urals

V. I. Ustinov and I. A. Baksheev p. 490 [abstract](#)

Model Equation of State for the Numerical Solution of the Problems of Multicomponent Filtration with Phase Transitions

A. V. Koldoba, E. V. Koldoba, and S. V. Skachkov p. 493 [abstract](#)

*Petrology*

Vol. 12, No. 3, 2004

Simultaneous English language translation of the journal is available from MAIK "Nauka / Interperiodica" (Russia).

*Petrology* ISSN 0869-5911.

Evolution of the Belomorian Belt: NORDSIM U–Pb Zircon Dating of the Chupa Paragneisses, Magmatism, and Metamorphic Stages

E. V. Bibikova, S. V. Bogdanova, V. A. Glebovitsky, S. Claesson, and T. Skiöld p. 195 [abstract](#)

Late Archean Granitoids of the Dambukinskii Block of the Dzhugdzhur-Stanovoy Fold Belt:

Formation and Transformation of the Continental Crust in the Early Precambrian

A. M. Larin, E. B. Sal'nikova, A. B. Kotov, V. A. Glebovitsky, V. P. Kovach, N. G. Berezhnaya, S. Z. Yakovleva, and M. D. Tolkachev p. 211 [abstract](#)

Structural and Compositional Characteristics of the Oldest Volcanic Glass in the Early Paleoproterozoic Boninite-Like Lavas of Southern Karelia

E. V. Sharkov, N. V. Trubkin, I. S. Krassivskaya, O. A. Bogatkov, A. V. Mokhov, A. V. Chistyakov, and K. A. Evseeva p. 227 [abstract](#)

Rhythmic Layering of the Kivakka Type: Geology, Petrography, Petrochemistry, and a Hypothesis for Its Formation

Ya. V. Bychkova and E. V. Koptev-Dvornikov p. 244 [abstract](#)

Distribution of Fe-Ni-Cu Sulfide Mineralization in the Rocks of the Burakovsko-Aganozerskii Layered Intrusion

V. S. Semenov, S. I. Korneev, O. A. Yakovleva, S. V. Semenov, A. B. Kol'tsov, N. G. Grinevich, and A. Kh. Zil'bershtein p. 265 [abstract](#)

Granitization of Amphibolites: 2. Characterization of Physical and Chemical Phenomena Related to Fluid Filtration through a Rock

L. I. Khodorevskaya p. 282 [abstract](#)

*Lithology and Mineral Resources*

Vol. 39, No. 3, 2004

Simultaneous English language translation of the journal is available from MAIK "Nauka / Interperiodica" (Russia).

Distributed worldwide by Kluwer Academic/Plenum Publishers. *Lithology and Mineral Resources* ISSN 0024-4902.

Glauconite at Different Stages of Lithogenesis in Lower Cambrian Rocks of Western Lithuania

T. A. Ivanovskaya and A. R. Geptner p. 191 [abstract](#)

Lithofacies Features and Formation Conditions of Aptian–Albian Rocks in the Western Ciscaucasus

A. E. Khardikov and I. A. Kholodnaya p. 203 [abstract](#)

Upper Oligocene Sediments of the Ciscaucasus, Volga–Don, and Mangyshlak Regions (Central Eastern Paratethys): Communication 1. Main Compositional and Structural Features

A. S. Stolyarov and E. I. Ivleva p. 213 [abstract](#)

Geochemistry of Calcareous–Silicate Rocks of the Kharagol Formation, Southern Baikal Region

L. Z. Reznitsky, S. I. Shkol'nik, and V. I. Levitsky p. 230 [abstract](#)

Carbon and Oxygen Isotopic Compositions of Carbonates from Precambrian Apatite-Bearing Carbonate Rocks of the Aldan Shield

V. N. Guliy and H. Wada p. 243 [abstract](#)

Saline Coals of the Ukraine

A. V. Ivanova p. 254 [abstract](#)

Holocene Siliceous Sediments in the Sea of Okhotsk

Yu. G. Volokhin, A. S. Astakhov, and N. G. Vashchenkova p. 259 [abstract](#)

REVIEWES

The Classification of Calcium Phosphates of Phosphorites

Yu. N. Zanin p. 281 [abstract](#)

Review of the Book by A.V. Maslov, M.T. Krupenin, E.Z. Gareev, and L.V. Anfimov,  
*The Riphean of the Western Slope of the Southern Urals*  
(Yekaterinburg: Zavarit. Inst. Geol. Geokhim. Ural. Otd. Ross. Akad. Nauk,  
2001, vol. 1, 351 p.; vol. 2, 134 p.; vol. 3, 130 p.; vol. 4, 103 p.)

V. G. Kuznetsov p. 283 [abstract](#)

Chronicle

The 75th Birthday of Petr Vasil'evich Zaritskii p. 284

Scientific Readings in Commemoration of Corresponding Member  
of the USSR Academy of Sciences Prof. Leonid Vasil'evich Pustovalov p. 286

---

***The people interested in receiving our newsletter through mail, can write to [acfonseca@geobrasil.net](mailto:acfonseca@geobrasil.net) or [revistadegeologia@yahoo.com.br](mailto:revistadegeologia@yahoo.com.br)***

***\*\*\*Le persone interessate in ricevere la nostra newsletter tramite e-mail, possono scrivere ad [acfonseca@geobrasil.net](mailto:acfonseca@geobrasil.net) ou [revistadegeologia@yahoo.com.br](mailto:revistadegeologia@yahoo.com.br).***